



# SAPPHIRE Grain



# Rat & Mouse KILLER

A SPECIALLY FORMULATED BAIT PROVEN TO  
CONTROL THE TOUGHEST RODENT INFESTATIONS

✓ PROFESSIONAL USE

✓ SINGLE FEED BAIT

0.005%  
BRODIFACOUM

100% RECYCLED  
BUCKET

5KG

E5510'a

## SAPPHIRE GRAIN

### A ready-to-use grain bait containing brodifacoum (0.005% w/w)

Authorisation n°: GB-2015-0904 / NI-2015-0904  
Approval holder : (GB) : LODI UK - Pensnett Trading Estate 3rd Avenue -KINGSWINFORD - West Midlands DY6 7FD United Kingdom- Tel. 0044 (0)1384 40 42 42 / (NI) : LODI SAS - PA des quatre routes 35390 Grand-Fougeray FRANCE.

- For use as a rodenticide by **Professional users with demonstrated competence** (trained professional) for the control of rats (*Rattus norvegicus* & *Rattus rattus*) and mice (*Mus musculus*) indoors and outdoors around buildings.

#### Use-specific instructions for use:

Application method(s) : Ready-to-use bait to be used in tamper-resistant bait stations or covered baiting points. Tamper-resistant bait stations must be:

- Strong enough to prevent entry or destruction by dogs.
- Strong enough to prevent entry or destruction by children under six years of age using hands, feet or objects reasonably expected to be available in the use area (e.g. household objects, toys, sticks, stones etc)
- Lockable or sealable so that children and dogs cannot gain access through the opening or mechanisms used to fill the bait compartments.
- Resistant to destruction or weakening from exposure to typical non-catastrophic weather (e.g. direct sunlight, extremes of temperature and humidity, rain, snow etc)

**Application rate:** For rats : 50-100 g of bait per bait station or covered and protected baiting points. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. For mice : 20-30 g of bait per bait station or covered and protected baiting points. If more than one bait station is needed, the minimum distance between bait stations should be of 2 meters.

Regularly check bait consumption and replace consumed or spoiled bait until consumption has stopped.

**When used in burrows:** Baits must be placed to minimise the exposure to non-target species and children.

Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled. **If used for pulsed baiting:** Replace eaten bait only after 3 days and then at maximum 7 day intervals. Collect any spilled bait and dead rodents. - Protect bait from the atmospheric conditions. Place the baiting points in areas non-labile to flooding. - Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt. - For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species - Baits must be placed to minimise the exposure to non-target species and children. Remove the remaining product at the end of treatment period. Repeat treatment in situations where there is evidence of new infestation. Follow any additional instructions provided by the CRRU UK Code of Best Practice (or equivalent). - Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it. - Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation. - Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve. - The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

-The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.). - Where possible, bait stations must be fixed to the ground or other structures. - Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened. - Bait should be secured so that it cannot be dragged away from the bait station. - Place the product out of the reach of children, birds, pets and farm animals and other non-target animals. -Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these. - Wear protective chemical resistant gloves during product handling phase (Nitril, EN374).- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product. - The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the CRRU UK Code of Best Practice (or equivalent). -Sites where burrow baiting is used should be visited more frequently than those where secure bait boxes are employed. - If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation. - If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure. Loose grain: Place the bait in the baiting point by using a dosage devise. Use a method to minimise dust (e.g. wet wiping).

**Batch** GR0541121

**Exp** 11/2023



Version 30-08-2024

SAPPHIRE GRAIN - brodifacoum (CAS 56073-10-0) : 0.005%

Danger H360D May damage the unborn child. H373 May cause damage to organs (blood) through prolonged or repeated exposure.

P201 Obtain special instructions before use. P202 Do not handle until all safety precautions have been read and understood. P260 Do not breathe dust. P280 Wear protective gloves. P308+P313 IF exposed or concerned: Get medical advice. P314 Get medical advice if you feel unwell. P405 Store locked up. P501 Dispose of contents/container in accordance with applicable regulations.

**LODI UK - Pensnett Trading Estate 3rd Avenue -KINGSWINFORD - West Midlands DY6 7FD - United Kingdom - Tel. 0044 (0)1384 40 42 42**



#### Use-specific risk mitigation measures:

Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign. -Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion. - To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. - Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities. -Do not use in areas where resistance to the active substance can be suspected. - Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment - Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant. - Do not wash the bait stations or utensils used in covered and protected bait points with water between applications. Dispose dead rodents in accordance with local requirements - In the UK poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>). To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements. Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance. When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

**Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment:** When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine. Antidote: Vitamin K1 administered by medical/veterinary personnel only. In case of:- Dermal exposure, wash skin with water and then with water and soap. - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes. - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet. **Bait stations must be labelled with the following information:** "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of accident, call a poison centre." UK medical professionals should contact the National Poisons Information Service ([www.npis.org](http://www.npis.org)) for further advice. Hazardous to wildlife.

#### Instructions for safe disposal of the product and its packaging and dead rodents:

At the end of the treatment, dispose of uneaten bait and the packaging in accordance with local requirements. Use of gloves is recommended. UK only: For information on disposal in the UK contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

#### Conditions of storage and shelf-life of the product under normal conditions of storage:

Shelf-life: 24 months - Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight. Store in places prevented from the access of children, birds, pets and farm animals. Keep only in original container.- Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait. Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them. This product contains a bittering agent and a dye

UFI : S68P-42SC-R00F-TCGH.

Marketed by: LODI UK  
Pensnett Trading Estate 3rd Avenue -  
KINGSWINFORD - West Midlands DY6 7FD -  
United Kingdom  
Tel. 0044 (0)1384 40 42 42

5Kg

  
LodiUK



**SAPPHIRE  
Grain**

**10KG**



**0.005%  
BRODIFACOUM**

**Rat & Mouse  
KILLER**



**A SPECIALLY FORMULATED BAIT PROVEN TO  
CONTROL THE TOUGHEST RODENT INFESTATIONS**

 **PROFESSIONAL USE**

 **SINGLE FEED BAIT**

E5511'a

## SAPPHIRE GRAIN

Version 11-10-2023

### A ready-to-use grain bait containing brodifacoum (0.005% w/w)

Authorisation n°: **GB-2015-0904 / NI-2015-0904** Approval holder : LODI UK - Pensnett Trading Estate 3rd Avenue -KINGSWINFORD - West Midlands DY6 7FD - United Kingdom / LODI SAS - PA des Quatre Routes 35390 Grand-Fougeray FRANCE  
For use as a rodenticide by **Professional users with demonstrated competence** (trained professional) for the control of rats (*Rattus norvegicus* & *Rattus rattus*) and mice (*Mus musculus*) indoors and outdoors around buildings.

#### Use-specific instructions for use:

Ready-to-use bait to be used in tamper-resistant bait stations, covered and protected baiting points or direct application of ready-to use bait into the burrow.

Tamper-resistant bait stations must be:

- Strong enough to prevent entry or destruction by dogs.
- Strong enough to prevent entry or destruction by children under six years of age using hands, feet or objects reasonably expected to be available in the use area (e.g. household objects, toys, sticks, stones, etc)
- Lockable or sealable so that children and dogs cannot gain access through the opening or mechanisms used to fill the bait compartments.
- Resistant to destruction or weakening from exposure to typical non-catastrophic weather (e.g. direct sunlight, extremes of temperature and humidity, rain, snow etc)

**Application rate:** For rats : 50-100 g of bait per bait station or covered and protected baiting points. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. For mice : 20-30 g of bait per bait station or covered and protected baiting points. If more than one bait station is needed, the minimum distance between bait stations should be of 2 meters.

Regularly check bait consumption and replace consumed or spoiled bait until consumption has stopped.

**When used in burrows:** Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled.

**If used for pulsed baiting:** Replace eaten bait only after 3 days and then at maximum 7 day intervals. Collect any spilled bait and dead rodents.

- Protect bait from the atmospheric conditions. Place the baiting points in areas non-labile to flooding. - Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt. - For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species - Baits must be placed to minimise the exposure to non-target species and children. Remove the remaining product at the end of treatment period. Repeat treatment in situations where there is evidence of new infestation. Follow any additional instructions provided by the CRRU UK Code of Best Practice (or equivalent).
- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it. - Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation. - Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve. - The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.). - Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened. - Bait should be secured so that it cannot be dragged away from the bait station. - Place the product out of the reach of children, birds, pets and farm animals and other non-target animals. -Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these. - Wear protective chemical resistant gloves during product handling phase (Nitril, EN374).- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product. - The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the CRRU UK Code of Best Practice (or equivalent).
- Sites where burrow baiting is used should be visited more frequently than those where secure bait boxes are employed. - If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation. - If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Loose grain: Place the bait in the baiting point by using a dosage devise. Use a method to minimise dust (e.g. wet wiping).

#### **Marketed by: LODI UK**

Pensnett Trading Estate 3rd Avenue -KINGSWINFORD - West Midlands  
DY6 7FD - United Kingdom - Tel. 0044 (0)1384 40 42 42



**10kg**

#### **SAPPHIRE GRAIN - brodifacoum (CAS 56073-10-0) : 0.005%**

Danger

**H360D May damage the unborn child. H373 May cause damage to organs (blood) through prolonged or repeated exposure.** P201 Obtain special instructions

before use. P202 Do not handle until all safety precautions have been read and understood. P260 Do not breathe dust. P280 Wear protective gloves.

P308+P313 IF exposed or concerned: Get medical advice. P314 Get medical advice if you feel unwell. P405 Store locked up. P501 Dispose of contents/container in accordance with applicable regulations.

**LODI UK - Pensnett Trading Estate 3rd Avenue -KINGSWINFORD - West Midlands DY6 7FD - United Kingdom**



#### Use-specific risk mitigation measures:

-Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign. -Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion. - To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice. - Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

-Do not use in areas where resistance to the active substance can be suspected. - Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment - Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant. - Do not wash the bait stations or utensils used in covered and protected bait points with water between applications. - Dispose dead rodents in accordance with local requirements - In the UK poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>). To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements. Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance. When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

#### Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

- Antidote: Vitamin K1 administered by medical/veterinary personnel only. - In case of:- Dermal exposure, wash skin with water and then with water and soap. - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes. - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet.

**Bait stations must be labelled with the following information:** "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of accident, call a poison centre." UK medical professionals should contact the National Poisons Information Service ([www.npis.org](http://www.npis.org)) for further advice. - Hazardous to wildlife.

#### Instructions for safe disposal of the product and its packaging and dead rodents:

At the end of the treatment, dispose of uneaten bait and the packaging in accordance with local requirements. Use of gloves is recommended. UK only: For information on disposal in the UK contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

#### Conditions of storage and shelf-life of the product under normal conditions of storage:

Shelf-life: 24 months - Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight. Store in places prevented from the access of children, birds, pets and farm animals. - Keep only in original container.- Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait. - Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them. - This product contains a bittering agent and a dye.

Batch GRO190224

Exp 02/2026

UFI : S68P-42SC-R00F-TCGH